

The transformation function analogous to (1.7.32) is

$$\langle \mathbf{x}' | \mathbf{p}' \rangle = \left[\frac{1}{(2\pi\hbar)^{3/2}} \right] \exp\left(\frac{i\mathbf{p}' \cdot \mathbf{x}'}{\hbar}\right), \quad (1.7.50)$$

so that

$$\psi_\alpha(\mathbf{x}') = \left[\frac{1}{(2\pi\hbar)^{3/2}} \right] \int d^3p' \exp\left(\frac{i\mathbf{p}' \cdot \mathbf{x}'}{\hbar}\right) \phi_\alpha(\mathbf{p}') \quad (1.7.51a)$$

and

$$\phi_\alpha(\mathbf{p}') = \left[\frac{1}{(2\pi\hbar)^{3/2}} \right] \int d^3x' \exp\left(\frac{-i\mathbf{p}' \cdot \mathbf{x}'}{\hbar}\right) \psi_\alpha(\mathbf{x}'). \quad (1.7.51b)$$

It is interesting to check the dimension of the wave functions. In one-dimensional problems the normalization requirement (1.6.8) implies that $|\langle x' | \alpha \rangle|^2$ has the dimension of inverse length, so the wave function itself must have the dimension of $(\text{length})^{-1/2}$. In contrast, the wave function in three-dimensional problems must have the dimension of $(\text{length})^{-3/2}$ because $|\langle \mathbf{x}' | \alpha \rangle|^2$ integrated over all spatial volume must be unity (dimensionless).

Problems

1. Prove

$$[AB, CD] = -AC\{D, B\} + A\{C, B\}D - C\{D, A\}B + \{C, A\}DB.$$

2. Suppose a 2×2 matrix X (not necessarily Hermitian, nor unitary) is written as

$$X = a_0 + \boldsymbol{\sigma} \cdot \mathbf{a},$$

where a_0 and $a_{1,2,3}$ are numbers.

a. How are a_0 and a_k ($k=1,2,3$) related to $\text{tr}(X)$ and $\text{tr}(\boldsymbol{\sigma}_k X)$?

b. Obtain a_0 and a_k in terms of the matrix elements X_{ij} .

3. Show that the determinant of a 2×2 matrix $\boldsymbol{\sigma} \cdot \mathbf{a}$ is invariant under

$$\boldsymbol{\sigma} \cdot \mathbf{a} \rightarrow \boldsymbol{\sigma} \cdot \mathbf{a}' \equiv \exp\left(\frac{i\boldsymbol{\sigma} \cdot \hat{\mathbf{n}}\phi}{2}\right) \boldsymbol{\sigma} \cdot \mathbf{a} \exp\left(\frac{-i\boldsymbol{\sigma} \cdot \hat{\mathbf{n}}\phi}{2}\right).$$

Find a'_k in terms of a_k when $\hat{\mathbf{n}}$ is in the positive z -direction and interpret your result.

4. Using the rules of bra-ket algebra, prove or evaluate the following:

a. $\text{tr}(XY) = \text{tr}(YX)$, where X and Y are operators;

b. $(XY)^\dagger = Y^\dagger X^\dagger$, where X and Y are operators;

c. $\exp[if(A)] = ?$ in ket-bra form, where A is a Hermitian operator whose eigenvalues are known;

d. $\sum_a \psi_a^*(\mathbf{x}') \psi_a(\mathbf{x}'')$, where $\psi_a(\mathbf{x}') = \langle \mathbf{x}' | a' \rangle$.

5. a. Consider two kets $|\alpha\rangle$ and $|\beta\rangle$. Suppose $\langle a' | \alpha \rangle$, $\langle a'' | \alpha \rangle$, ... and $\langle a' | \beta \rangle$, $\langle a'' | \beta \rangle$, ... are all known, where $|a'\rangle$, $|a''\rangle$, ... form a complete set of base kets. Find the matrix representation of the operator $|\alpha\rangle\langle\beta|$ in that basis.

b. We now consider a spin $\frac{1}{2}$ system and let $|\alpha\rangle$ and $|\beta\rangle$ be $|s_z = \hbar/2\rangle$ and $|s_x = \hbar/2\rangle$, respectively. Write down explicitly the square matrix that corresponds to $|\alpha\rangle\langle\beta|$ in the usual (s_z diagonal) basis.

6. Suppose $|i\rangle$ and $|j\rangle$ are eigenkets of some Hermitian operator A . Under what condition can we conclude that $|i\rangle + |j\rangle$ is also an eigenket of A ? Justify your answer.

7. Consider a ket space spanned by the eigenkets $\{|a'\rangle\}$ of a Hermitian operator A . There is no degeneracy.

a. Prove that

$$\prod_{a'} (A - a')$$

is the null operator.

b. What is the significance of

$$\prod_{a'' \neq a'} \frac{(A - a'')}{(a' - a'')}?$$

c. Illustrate (a) and (b) using A set equal to S_z of a spin $\frac{1}{2}$ system.

8. Using the orthonormality of $|+\rangle$ and $|-\rangle$, prove

$$[S_i, S_j] = i\epsilon_{ijk}\hbar S_k, \quad \{S_i, S_j\} = \left(\frac{\hbar^2}{2}\right)\delta_{ij},$$

where

$$S_x = \frac{\hbar}{2}(|+\rangle\langle-| + |-\rangle\langle+|), \quad S_y = \frac{i\hbar}{2}(-|+\rangle\langle-| + |-\rangle\langle+|),$$

$$S_z = \frac{\hbar}{2}(|+\rangle\langle+| - |-\rangle\langle-|).$$


9. Construct $|\mathbf{S} \cdot \hat{\mathbf{n}}; +\rangle$ such that

$$\mathbf{S} \cdot \hat{\mathbf{n}} |\mathbf{S} \cdot \hat{\mathbf{n}}; +\rangle = \left(\frac{\hbar}{2}\right) |\mathbf{S} \cdot \hat{\mathbf{n}}; +\rangle$$

where $\hat{\mathbf{n}}$ is characterized by the angles shown in the figure. Express your answer as a linear combination of $|+\rangle$ and $|-\rangle$. [Note: The answer is

$$\cos\left(\frac{\beta}{2}\right)|+\rangle + \sin\left(\frac{\beta}{2}\right)e^{i\alpha}|-\rangle.$$

But do not just verify that this answer satisfies the above eigenvalue equation. Rather, treat the problem as a straightforward eigenvalue



problem. Also do not use rotation operators, which we will introduce later in this book.]

10. The Hamiltonian operator for a two-state system is given by

$$H = a(|1\rangle\langle 1| - |2\rangle\langle 2| + |1\rangle\langle 2| + |2\rangle\langle 1|),$$

where a is a number with the dimension of energy. Find the energy eigenvalues and the corresponding energy eigenkets (as linear combinations of $|1\rangle$ and $|2\rangle$).

11. A two-state system is characterized by the Hamiltonian

$$H = H_{11}|1\rangle\langle 1| + H_{22}|2\rangle\langle 2| + H_{12}[|1\rangle\langle 2| + |2\rangle\langle 1|]$$

where H_{11} , H_{22} , and H_{12} are real numbers with the dimension of energy, and $|1\rangle$ and $|2\rangle$ are eigenkets of some observable ($\neq H$). Find the energy eigenkets and corresponding energy eigenvalues. Make sure that your answer makes good sense for $H_{12} = 0$. (You need not solve this problem from scratch. The following fact may be used without proof:

$$(\mathbf{S} \cdot \hat{\mathbf{n}})|\hat{\mathbf{n}}; +\rangle = \frac{\hbar}{2}|\hat{\mathbf{n}}; +\rangle,$$

with $|\hat{\mathbf{n}}; +\rangle$ given by

$$|\hat{\mathbf{n}}; +\rangle = \cos \frac{\beta}{2}|+\rangle + e^{i\alpha} \sin \frac{\beta}{2}|-\rangle,$$

where β and α are the polar and azimuthal angles, respectively, that characterize $\hat{\mathbf{n}}$.

12. A spin $\frac{1}{2}$ system is known to be in an eigenstate of $\mathbf{S} \cdot \hat{\mathbf{n}}$ with eigenvalue $\hbar/2$, where $\hat{\mathbf{n}}$ is a unit vector lying in the xz -plane that makes an angle γ with the positive z -axis.

- a. Suppose S_x is measured. What is the probability of getting $+\hbar/2$?
b. Evaluate the dispersion in S_x , that is,

$$\langle (S_x - \langle S_x \rangle)^2 \rangle.$$

(For your own peace of mind check your answers for the special cases $\gamma = 0, \pi/2$, and π .)

13. A beam of spin $\frac{1}{2}$ atoms goes through a series of Stern-Gerlach-type measurements as follows:

- a. The first measurement accepts $s_z = \hbar/2$ atoms and rejects $s_z = -\hbar/2$ atoms.
b. The second measurement accepts $s_n = \hbar/2$ atoms and rejects $s_n = -\hbar/2$ atoms, where s_n is the eigenvalue of the operator $\mathbf{S} \cdot \hat{\mathbf{n}}$, with $\hat{\mathbf{n}}$ making an angle β in the xz -plane with respect to the z -axis.
c. The third measurement accepts $s_z = -\hbar/2$ atoms and rejects $s_z = \hbar/2$ atoms.

What is the intensity of the final $s_z = -\hbar/2$ beam when the $s_z = \hbar/2$ beam surviving the first measurement is normalized to unity? How must we orient the second measuring apparatus if we are to maximize the intensity of the final $s_z = -\hbar/2$ beam?

14. A certain observable in quantum mechanics has a 3×3 matrix representation as follows:

$$\frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}.$$

- a. Find the normalized eigenvectors of this observable and the corresponding eigenvalues. Is there any degeneracy?
b. Give a physical example where all this is relevant.
15. Let A and B be observables. Suppose the simultaneous eigenkets of A and B $\{|a', b'\rangle\}$ form a *complete* orthonormal set of base kets. Can we always conclude that

$$[A, B] = 0?$$

If your answer is yes, prove the assertion. If your answer is no, give a counterexample.

16. Two Hermitian operators anticommute:

$$\{A, B\} = AB + BA = 0.$$

Is it possible to have a simultaneous (that is, common) eigenket of A and B ? Prove or illustrate your assertion.

17. Two observables A_1 and A_2 , which do not involve time explicitly, are known not to commute,

$$[A_1, A_2] \neq 0,$$

yet we also know that A_1 and A_2 both commute with the Hamiltonian:

$$[A_1, H] = 0, \quad [A_2, H] = 0.$$

Prove that the energy eigenstates are, in general, degenerate. Are there exceptions? As an example, you may think of the central-force problem $H = \mathbf{p}^2/2m + V(r)$, with $A_1 \rightarrow L_z$, $A_2 \rightarrow L_x$.

18. a. The simplest way to derive the Schwarz inequality goes as follows. First, observe

$$(\langle \alpha | + \lambda^* \langle \beta |) \cdot (|\alpha\rangle + \lambda |\beta\rangle) \geq 0$$

for any complex number λ ; then choose λ in such a way that the preceding inequality reduces to the Schwarz inequality.

- b. Show that the equality sign in the generalized uncertainty relation holds if the state in question satisfies

$$\Delta A |\alpha\rangle = \lambda \Delta B |\alpha\rangle$$

with λ purely *imaginary*.

- c. Explicit calculations using the usual rules of wave mechanics show that the wave function for a Gaussian wave packet given by

$$\langle x' | \alpha \rangle = (2\pi d^2)^{-1/4} \exp \left[\frac{i \langle p \rangle x'}{\hbar} - \frac{(x' - \langle x \rangle)^2}{4d^2} \right]$$

satisfies the minimum uncertainty relation

$$\sqrt{\langle (\Delta x)^2 \rangle} \sqrt{\langle (\Delta p)^2 \rangle} = \frac{\hbar}{2}.$$

Prove that the requirement

$$\langle x' | \Delta x | \alpha \rangle = (\text{imaginary number}) \langle x' | \Delta p | \alpha \rangle$$

is indeed satisfied for such a Gaussian wave packet, in agreement with (b).

19. a. Compute

$$\langle (\Delta S_x)^2 \rangle \equiv \langle S_x^2 \rangle - \langle S_x \rangle^2,$$

where the expectation value is taken for the $S_z +$ state. Using your result, check the generalized uncertainty relation

$$\langle (\Delta A)^2 \rangle \langle (\Delta B)^2 \rangle \geq \frac{1}{4} |\langle [A, B] \rangle|^2,$$

with $A \rightarrow S_x$, $B \rightarrow S_y$.

- b. Check the uncertainty relation with $A \rightarrow S_x$, $B \rightarrow S_y$ for the $S_x +$ state.
20. Find the linear combination of $|+\rangle$ and $|-\rangle$ kets that maximizes the

uncertainty product

$$\langle (\Delta S_x)^2 \rangle \langle (\Delta S_y)^2 \rangle.$$

Verify explicitly that for the linear combination you found, the uncertainty relation for S_x and S_y is not violated.

21. Evaluate the x - p uncertainty product $\langle (\Delta x)^2 \rangle \langle (\Delta p)^2 \rangle$ for a one-dimensional particle confined between two rigid walls

$$V = \begin{cases} 0 & \text{for } 0 < x < a, \\ \infty & \text{otherwise.} \end{cases}$$

Do this for both the ground and excited states.

22. Estimate the rough order of magnitude of the length of time that an ice pick can be balanced on its point if the only limitation is that set by the Heisenberg uncertainty principle. Assume that the point is sharp and that the point and the surface on which it rests are hard. You may make approximations which do not alter the general order of magnitude of the result. Assume reasonable values for the dimensions and weight of the ice pick. Obtain an approximate numerical result and express it in *seconds*.
23. Consider a three-dimensional ket space. If a certain set of orthonormal kets—say, $|1\rangle$, $|2\rangle$, and $|3\rangle$ —are used as the base kets, the operators A and B are represented by

$$A \doteq \begin{pmatrix} a & 0 & 0 \\ 0 & -a & 0 \\ 0 & 0 & -a \end{pmatrix}, \quad B \doteq \begin{pmatrix} b & 0 & 0 \\ 0 & 0 & -ib \\ 0 & ib & 0 \end{pmatrix}$$

with a and b both real.

- a. Obviously A exhibits a degenerate spectrum. Does B also exhibit a degenerate spectrum?
- b. Show that A and B commute.
- c. Find a new set of orthonormal kets which are simultaneous eigenkets of both A and B . Specify the eigenvalues of A and B for each of the three eigenkets. Does your specification of eigenvalues completely characterize each eigenket?
24. a. Prove that $(1/\sqrt{2})(1 + i\sigma_x)$ acting on a two-component spinor can be regarded as the matrix representation of the rotation operator about the x -axis by angle $-\pi/2$. (The minus sign signifies that the rotation is clockwise.)
- b. Construct the matrix representation of S_z when the eigenkets of S_y are used as base vectors.
25. Some authors define an *operator* to be real when every member of its matrix elements $\langle b' | A | b'' \rangle$ is real in some representation ($\{|b'\rangle\}$ basis in this case). Is this concept representation independent, that is, do the

- matrix elements remain real even if some basis other than $\{|b'\rangle\}$ is used? Check your assertion using familiar operators such as S_y and S_z (see Problem 24) or x and p_x .
26. Construct the transformation matrix that connects the S_z diagonal basis to the S_x diagonal basis. Show that your result is consistent with the general relation

$$U = \sum_r |b^{(r)}\rangle \langle a^{(r)}|.$$

27. a. Suppose that $f(A)$ is a function of a Hermitian operator A with the property $A|a'\rangle = a'|a'\rangle$. Evaluate $\langle b''|f(A)|b'\rangle$ when the transformation matrix from the a' basis to the b' basis is known.
- b. Using the continuum analogue of the result obtained in (a), evaluate

$$\langle \mathbf{p}''|F(r)|\mathbf{p}'\rangle.$$

Simplify your expression as far as you can. Note that r is $\sqrt{x^2 + y^2 + z^2}$, where x , y , and z are operators.

28. a. Let x and p_x be the coordinate and linear momentum in one dimension. Evaluate the classical Poisson bracket

$$[x, F(p_x)]_{\text{classical}}.$$

- b. Let x and p_x be the corresponding quantum-mechanical operators this time. Evaluate the commutator

$$\left[x, \exp\left(\frac{ip_x a}{\hbar}\right) \right].$$

- c. Using the result obtained in (b), prove that

$$\exp\left(\frac{ip_x a}{\hbar}\right)|x'\rangle, \quad (x|x'\rangle = x'|x'\rangle)$$

is an eigenstate of the coordinate operator x . What is the corresponding eigenvalue?

29. a. On page 247, Gottfried (1966) states that

$$[x_i, G(\mathbf{p})] = i\hbar \frac{\partial G}{\partial p_i}, \quad [p_i, F(\mathbf{x})] = -i\hbar \frac{\partial F}{\partial x_i}$$

can be “easily derived” from the fundamental commutation relations for all functions of F and G that can be expressed as power series in their arguments. Verify this statement.

- b. Evaluate $[x^2, p^2]$. Compare your result with the classical Poisson bracket $[x^2, p^2]_{\text{classical}}$.

30. The translation operator for a finite (spatial) displacement is given by

$$\mathcal{T}(\mathbf{l}) = \exp\left(\frac{-i\mathbf{p}\cdot\mathbf{l}}{\hbar}\right),$$

where \mathbf{p} is the momentum operator.

- a. Evaluate

$$[x_i, \mathcal{T}(\mathbf{l})].$$

- b. Using (a) (or otherwise), demonstrate how the expectation value $\langle \mathbf{x} \rangle$ changes under translation.

31. In the main text we discussed the effect of $\mathcal{T}(d\mathbf{x}')$ on the position and momentum eigenkets and on a more general state ket $|\alpha\rangle$. We can also study the behavior of expectation values $\langle \mathbf{x} \rangle$ and $\langle \mathbf{p} \rangle$ under infinitesimal translation. Using (1.6.25), (1.6.45), and $|\alpha\rangle \rightarrow \mathcal{T}(d\mathbf{x}')|\alpha\rangle$ only, prove $\langle \mathbf{x} \rangle \rightarrow \langle \mathbf{x} \rangle + d\mathbf{x}'$, $\langle \mathbf{p} \rangle \rightarrow \langle \mathbf{p} \rangle$ under infinitesimal translation.

32. a. Verify (1.7.39a) and (1.7.39b) for the expectation value of p and p^2 from the Gaussian wave packet (1.7.35).

- b. Evaluate the expectation value of p and p^2 using the momentum-space wave function (1.7.42).

33. a. Prove the following:

$$(i) \quad \langle p'|x|\alpha\rangle = i\hbar \frac{\partial}{\partial p'} \langle p'|\alpha\rangle,$$

$$(ii) \quad \langle \beta|x|\alpha\rangle = \int dp' \phi_\beta^*(p') i\hbar \frac{\partial}{\partial p'} \phi_\alpha(p'),$$

where $\phi_\alpha(p') = \langle p'|\alpha\rangle$ and $\phi_\beta(p') = \langle p'|\beta\rangle$ are momentum-space wave functions.

- b. What is the physical significance of

$$\exp\left(\frac{ix\Xi}{\hbar}\right),$$

where x is the position operator and Ξ is some number with the dimension of momentum? Justify your answer.